arXiv:1403.4371v1 [math.AG] 18 Mar 2014

THE SIGNED EULER CHARACTERISTIC OF VERY AFFINE VARIETIES

NERO BUDUR AND BOTONG WANG

Abstract. A conjecture of J. Huh and B. Sturmfels predicts that the sign of the Euler characteristic of a complex very affine variety depends only on the parity of the dimension. The conjecture is true for locally complete intersections. Beyond this case, we construct counterexamples with arbitrarily bad failure.

1. Introduction

Let X be a closed irreducible subvariety of $(\mathbb{C}^*)^n$. In the literature, this is called a very affine variety. When X is a locally complete intersection, $(-1)^{\dim(X)}\chi(X) \geq 0$. This follows from generic vanishing results for perverse sheaves on $(\mathbb{C}^*)^n$ due to Loeser-Sabbah [LS] (see also Gabber-Loeser [GL]), together with the well-known fact that for a locally complete intersection the shifted sheaf $\mathbb{C}_X[\dim X]$ is perverse. For the smooth case, see also [H].

It was conjectured by Huh and Sturmfels [HS, page 6] that the same is true for any closed irreducible subvariety X of $(\mathbb{C}^*)^n$. In this note, we construct counterexamples by displaying singular surfaces in $(\mathbb{C}^*)^4$ with arbitrary negative Euler characteristics.

2. Construction

We start with a smooth surface U in $(\mathbb{C}^*)^4$ defined as

$$U = \{(w, x, y, z) \in (\mathbb{C}^*)^4 \mid w + y = x + z = 1\},$$

where w, x, y, z are the coordinates in $(\mathbb{C}^*)^4$.

We define an action of $\mathbb{Z}/n\mathbb{Z}$ on $(\mathbb{C}^*)^4$. Let $\xi \in \mathbb{C}$ be an n-th primitive root of unity. We set $\xi(w, x, y, z) = (\xi w, \xi x, \xi^{-1} y, \xi^{-1} z)$. This defines a $\mathbb{Z}/n\mathbb{Z}$ action on $(\mathbb{C}^*)^4$ by translations. Hence, the quotient $(\mathbb{C}^*)^4/(\mathbb{Z}/n\mathbb{Z})$ is again a commutative affine algebraic group. Such algebraic group has to be isomorphic to $(\mathbb{C}^*)^4$. In fact, we can give an explicit description of the quotient map, which we denote by $p_n : (\mathbb{C}^*)^4 \to (\mathbb{C}^*)^4$,

$$p_n : (w, x, y, z) \mapsto (w^n, \frac{w}{x}, wy, wz).$$

We denote the image $p_n(U)$ by U_n. Then U_n is an irreducible subvariety of $(\mathbb{C}^*)^n$.

Theorem 2.1. *When n is odd, $\chi(U_n) = \frac{3-n}{2}$.*

The proof of the theorem will be in the next section.

The first author was partly sponsored by the Simons Foundation and NSA.

3. Euler characteristic

We will see that U_n has only isolated singularities, which are analytically equivalent to the transverse intersection of two smooth surfaces in $\mathbb{C}^4$. Moreover, the normalisation of U_n will be isomorphic to U. This allows us to compute the Euler characteristic of U_n by counting the number of singular points on U_n. Throughout this section, we assume n is odd.

Lemma 3.1. *For any $1 \leq i \leq n-1$, $U \cap \xi^i U$ has exact one point, and the intersection is transverse. Furthermore, $U \cap \xi^i U \cap \xi^j U = 0$ for $1 \leq i < j \leq n-1$.*

Proof. Recall that $U = \{(w,x,y,z) \in (\mathbb{C}^*)^4 \mid w+y = x+z = 1\}$. Hence

$$\xi^i U = \{(w,x,y,z) \in (\mathbb{C}^*)^4 \mid \xi^{-i}w + \xi^i y = \xi^{-i}x + \xi^i z = 1\}.$$

A direct computation shows $U \cap \xi^i U = \{(\frac{\xi^i}{1+\xi^i}, \frac{1}{1+\xi^i}, \frac{\xi^i}{1+\xi^i}, \frac{1}{1+\xi^i})\}$. Since the intersection is defined by 4 linear equations, clearly it is transverse. The last part is obvious, since $\frac{\xi^i}{1+\xi^i} \neq \frac{\xi^j}{1+\xi^j}$ for $1 \leq i < j \leq n-1$. □

Corollary 3.2. *U_n has $\frac{n-1}{2}$ isolated singular points. Moreover, the germ of U_n at any singular point is analytically equivalent to the germ of $\{w = x = 0\} \cup \{y = z = 0\}$ in $\mathbb{C}^4$ at origin. In other words, locally the singularity is obtained by the transverse intersection of two smooth surfaces.*

Proof. The quotient map $p_n : (\mathbb{C}^*)^4 \to (\mathbb{C}^*)^4$ restricts to a map $q_n : U \to U_n$. The map q_n is an isomorphism on $U - \bigcup_{1 \leq i \leq n-1}(U \cap \xi^i U)$. Every intersection $U \cap \xi^i U$ will create a singular point on U_n, which is the image of $U \cap \xi^i U$ under q_n. Since the intersection $U \cap \xi^i U$ is transverse, and since this intersection is not contained in any other $\xi^j U$, the corresponding singular point in U_n is locally isomorphic to the transverse intersection of two smooth surfaces. Notice that $\xi^{-i}(U \cap \xi^i U) = U \cap \xi^{n-i} U$. $U \cap \xi^i U$ and $U \cap \xi^{n-i} U$ give the same singular point in U_n. On the other hand, each singular point in U_n comes from exactly two such intersections. Therefore, U_n has exactly $\frac{n-1}{2}$ singular points. □

Proof of Theorem 2.1. The map $\mathbb{C}^2 \to \mathbb{C}^4$, defined by $(a,b) \mapsto (a, b, 1-a, 1-b)$ induces an isomorphism between $(\mathbb{C} - \{0,1\}) \times (\mathbb{C} - \{0,1\})$ and U. Therefore, by Kunneth's formula $\chi(U) = 1$. According to the corollary, U_n is obtained from U by attaching $\frac{n-1}{2}$ pairs of points. Thus $\chi(U_n) = 1 - \frac{n-1}{2} = \frac{3-n}{2}$. □

Example 3.3. The smallest n for which U_n is a counterexample to the above-mentioned conjecture is $n = 5$: $\chi(U_5) = -1$. Using a computer algebra program reveals that the equations for U_5 are quite complicated. It is the common zero locus in $(\mathbb{C}^*)^4$ of the

following five equations in 4 variables:

$$
\begin{aligned}
0 &= t_2^2t_4^2 + t_2^2t_3 - t_2^2t_4 - 2t_2t_3t_4 - t_2t_3 + {t_3}^2 + t_2t_4,\\
0 &= t_2{t_3}^2t_4 + 2t_2{t_3}^2 - {t_3}^3 - 3t_2t_3t_4 - t_1t_2 - t_2t_3 + {t_3}^2 + t_2t_4 + t_1,\\
0 &= {t_3}^4 - {t_3}^3t_4 + t_1t_2{t_4}^2 - 5t_2t_3{t_4}^2 + t_1t_2t_3 - 5t_2{t_3}^2 + 3{t_3}^3 - t_1t_2t_4 - t_1t_3t_4 + 5t_2t_3t_4 +\\
&\quad + 2{t_3}^2t_4 + 2t_2{t_4}^2 - 3t_1t_3 + 2t_2t_3 - 2{t_3}^2 + 3t_1t_4 - 2t_2t_4,\\
0 &= t_2{t_3}^3 + 3t_2{t_3}^2 - {t_3}^3 + t_1t_2t_4 - 5t_2t_3t_4 - 3t_1t_2 - t_1t_3 - 2t_2t_3 + 2{t_3}^2 + 2t_2t_4 + 3t_1,\\
0 &= t_1t_2t_3{t_4}^2 - {t_3}^3{t_4}^2 + t_1t_2{t_4}^3 - 5t_2t_3{t_4}^3 + t_1t_2{t_3}^2 - t_1{t_3}^2t_4 + 3{t_3}^3t_4 - 4t_1t_2{t_4}^2\\
&\quad - t_1t_3{t_4}^2 + 7t_2t_3{t_4}^2 + 2{t_3}^2{t_4}^2 + 2t_2{t_4}^3 - 3t_1t_2t_3 + 2t_1{t_3}^2 + 12t_2{t_3}^2 - 6{t_3}^3 + 3t_1t_2t_4 +\\
&\quad + 3t_1t_3t_4 - 15t_2t_3t_4 - 3{t_3}^2t_4 + 3t_1{t_4}^2 - 3t_2{t_4}^2 - {t_1}^2 - 5t_1t_2 - t_1t_3 - 6t_2t_3 + 6{t_3}^2 -\\
&\quad - 4t_1t_4 + 6t_2t_4 + 6t_1.
\end{aligned}
$$

References

[GL] O. Gabber and F. Loeser, Faisceaux pervers l-adiques sur un tore. Duke Math. J. 83 (1996), no. 3, 501–606. 1

[H] J. Huh, The maximum likelihood degree of a very affine variety. Compos. Math. 149 (2013), no. 8, 1245–1266. 1

[HS] J. Huh and B. Sturmfels, Likelihood Geometry. arXiv:1305.7462. To appear in Combinatorial Algebraic Geometry, Lecture Notes in Mathematics 2108, Springer, 2014. 1

[LS] F. Loeser and C. Sabbah, Caractérisation des D-modules hypergéométriques irréductibles sur le tore. C. R. Acad. Sci. Paris Sér. I Math. 312 (1991), no. 10, 735–738. II, C. R. Acad. Sci. Paris Sér. I Math. 315 (1992), no. 12, 1263–1264. 1

E-mail address: Nero.Budur@wis.kuleuven.be

KU Leuven and University of Notre Dame

Current address: KU Leuven, Department of Mathematics, Celestijnenlaan 200B, B-3001 Leuven, Belgium

E-mail address: bwang3@nd.edu

University of Notre Dame

Current address: Department of Mathematics, 255 Hurley Hall, IN 46556, USA